

Dosage Recommendation



Cerebrolysin®

Reconnecting Neurons.
Empowering for Life.

Dosage recommendation

Disorder	Daily dosage	Initiation of treatment	Duration of treatment
Stroke	20 - 50 ml	as soon as possible	10 - 21 days
Traumatic brain injury	20 - 50 ml	as soon as possible	7 - 30 days
Vascular dementia	10 - 30 ml	as soon as possible	1 cycle: 5 days weekly/4 weeks 2-4 cycles per year
Alzheimer's disease	10 - 30 ml	as soon as possible	1 cycle: 5 days weekly/4 weeks 2-4 cycles per year

Route of administration




IV injection	up to 10ml	undiluted	inject slowly over 3 minutes
IM injection	up to 5ml	undiluted	inject slowly over 3 minutes
The IV or IM injection has to be administered immediately after opening the ampoule!			



IV infusion	10ml - 50ml	diluted to at least 100ml total volume with Saline, Ringer solution or 5% glucose solution	Infuse within 15 minutes
To guarantee sterility, always use disposable one-way infusion sets and cannulas! Start the infusion as quickly as possible after dilution! Cerebrolysin should not be mixed with balanced amino acid solutions, vitamins and cardiovascular medicinal products.			

Dilution procedures for IV infusion



Flush with sodium chloride solution before & after infusion

Safe administration – sterility aspects

Special precautions to guarantee sterility must be taken during dilution and administration of Cerebrolysin:

- Remove solution from ampoules immediately before use
- Do not leave an open ampoule on the treatment table
- Always use only disposable one-way IV infusions sets and cannulas
- When Cerebrolysin is administered via a long-term IV catheter, the catheter has to be rinsed before and after the application with physiological sodium chloride solution
- Pay special attention to recommended infusion/injection times on previous page

Conditions for storage

- Cerebrolysin must be stored at room temperature not exceeding 25°C
- Do **not** freeze!
- Has to be protected from light (in the carton)

ABBREVIATED PRESCRIBING INFORMATION - Cerebrolysin. Name of the medicinal product: Cerebrolysin - Solution for injection. Qualitative and quantitative composition: One ml contains 215.2 mg of Cerebrolysin concentrate in aqueous solution. List of excipients: Sodium hydroxide and water for injection. Therapeutic indications: For treatment of cerebrovascular disorders. Especially in the following indications: Senile dementia of Alzheimer's type. Vascular dementia. Stroke. Craniocerebral trauma (commotio and contusio). Contraindications: Hypersensitivity to one of the components of the drug, epilepsy, severe renal impairment. Marketing Authorisation Holder: EVER Neuro Pharma GmbH, A-4866 Unterach. Only available on prescription and in pharmacies. More information about pharmaceutical form, posology and method of administration, special warnings and precautions for use, interaction with other medicinal products and other forms of interaction, fertility, pregnancy and lactation, effects on ability to drive and use machines, undesirable effects, overdose, pharmacodynamics properties, pharmacokinetic properties, preclinical safety data, incompatibilities, shelf life, special precautions for storage, nature and contents of the container and special precautions for disposal is available in the summary of product characteristics.

Copyright © 2016 by EVER Neuro Pharma GmbH, Oberburgau 3, 4866 Unterach, Austria. All rights reserved. No part of this brochure may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without permission in writing from the publisher. Cerebrolysin is a registered trademark of EVER Neuro Pharma GmbH, 4866 Unterach, Austria