



STROKE

TBI

DEMENTIA

## Early treatment effects after TBI

### CAPTAIN – Safety and Efficacy of Cerebrolysin® in Neurorecovery after moderate-severe Traumatic Brain Injury

Muresanu, Dafin F., et al., „Efficacy and safety of cerebrolysin in neurorecovery after moderate-severe traumatic brain injury: results from the CAPTAIN II trial.“ Neurological Sciences: 1-11

#### The effective treatment after TBI

- Early treatment effects
- Significantly reduced depression
- Improvement of memory and concentration


**Cerebrolysin®**

Reconnecting Neurons.  
Empowering for Life.

# Treat with Cerebrolysin® as soon as possible after TBI

## Beneficial effects already on day 10

- Early independence
- Patients are less dependent on caregivers early on
- High potential for earlier discharge



**Title:** Efficacy and safety of Cerebrolysin® in Neurorecovery after moderate-severe traumatic brain injury:  
Results from the CAPTAIN II trial

Patients: 80 Cerebrolysin® group + 59 Placebo group = 139 patients in total

**Treatment:** Treatment Cycle 1 = Day 1 – 10 : 50 ml/day  
Treatment Cycle 2 = Day 31 – 40 : 10 ml/day  
Treatment Cycle 3 = Day 61 – 70 : 10 ml/day